


1 / 14

FIG. 1

APPROVED	O.G. FIG.
BY	CLASS
DRAFTSMAN	SUBCLASS



M 1 2 3


SDS-PAGE analysis of PEG-modified hA5B7 Fab'

Samples of unmodified hA5B7 Fab' (lane 1), hinge-modified Fab' (lane 2), and randomly-modified Fab' (lane 3) were prepared with non-reducing sample buffer, and 1.5µg of each loaded onto a 4-20% gradient Tris-glycine gel. Standard protein markers (lane M) were also run. These comprised myosin (200kDa), beta-galactosidase (116.3kDa), phosphorylase b (97.4 kDa), bovine serum albumin (66.3kDa), glutamate dehydrogenase (55.4kDa), lactate dehydrogenase (36.5 kDa), carbonic anhydrase (31kDa), trypsin inhibitor (21.5kDa), lysozyme (14.4kDa), aprotinin (6kDa) and insulin B & A chains (3.5 & 2.5kDa).

Following electrophoresis, the gel was stained with coomassie blue.


2/14

FIG. 2

Pharmacokinetics of ¹²⁵I labelled hA5B7 Fab' in rats

3/14

FIG. 3



M 1 2 3

SDS-PAGE analysis of hTNF40 Fab'-PEG conjugates

Samples of hTNF40 Fab'-PEG (25kDa) prepared by random conjugation (lane 1), Fab'-PEG (25kDa) prepared by hinge attachment (lane 2), and Fab'-PEG (40kDa) prepared by hinge attachment (lane 3) were prepared with non-reducing sample buffer, and 1.5 μ g of each loaded onto a 4-20% gradient Tris-glycine gel. Standard protein markers (lane M) were also run. These comprised myosin (200kDa), beta-galactosidase (116.3kDa), phosphorylase b (97.4 kDa), bovine serum albumin (66.3kDa), glutamate dehydrogenase (55.4kDa), lactate dehydrogenase (36.5 kDa), carbonic anhydrase (31kDa), trypsin inhibitor (21.5kDa), lysozyme (14.4kDa), aprotinin (6kDa) and insulin B & A chains (3.5 & 2.5kDa).

Following electrophoresis, the gel was stained with coomassie blue.


APPROVED	O.G. FIG.
BY	CLASS
DRAFTSMAN	SUBCLASS

SECRETED

4 / 14

APPROVED	O.G. FIG.	
BY	CLASS	SUBCLASS
DRAFTSMAN		


FIG. 4 Comparison between different PEG Fab' 's, Fab' of hTNF40.4, and hTNF40.4 in the L929 assay



5/14

APPROVED	O.G. FIG.
BY	CLASS SUBCLASS
BRAFTSHAH	

FIG. 5 Pharmacokinetics of 125 I-labelled hTNF40 Fab'-PEG in rats



APPROVED	O.G. FIG.	
BY	CLASS	SUBCLASS
DRAFTSMAN		

RECEIVED


WO 98/25971

09/214251

PCT/GB97/03400

6/14


FIG. 6 Pharmacokinetics of ¹¹¹In labelled hTNF40 in rats



7/14

APPROVED	O.G. FIG.
BY	CLASS
DRAFTSMAN	SUBCLASS

FIG. 7




HPLC gel filtration of hTNF40 Fab', Fab'-PEG and Fab'(PEG)₂

DuPont Zorbax GF-250 column run at 1ml/min in 0.2M phosphate buffer pH7.0

APPROVED	O.G. FIG.	
BY	CLASS	SUBCLASS
DRAFTSMAN		


FIG. 8 Pharmacokinetics of 125-I labelled TN3 in rats



9/14

APPROVED	O.G. FIG.	
BY	CLASS	SUBCLASS
DRAFTSMAN		


FIG. 9 ECE15 : Pharmacokinetics of 125 I labelled TN3 in rats
 Fab' PEGylated via the hinge with 25K PEG



10/14

APPROVED	O.G. FIG.
BY	CLASS
DRAFTSMAN	SUBCLASS


FIG. 10 Pharmacokinetics of 125 I-labelled hTNF40 Fab'-PEG in rats



11/14

SDS-PAGE analysis under non reducing
conditions of Fab'-PEG
(5kDa) prepared using a vinylsulphone or
iodoacetamide reagent

APPROVED BY	O.G. FIG.	
	CLASS	SUBCLASS
DRAFTSMAN		




1. Molecular weight marker proteins
2. Purified Fab' (also containing F(ab')₂)
3. Fab'-PEG (5kDa, VS linker) reaction mix
4. Fab'-PEG (5kDa, IA linker) reaction mix
5. Fab'-PEG (5kDa, VS linker) reaction mix
6. Fab'-PEG (5kDa, IA linker) reaction mix

FIG. 11

FIG. 12

DAD1 A, Sig=280,4 Ref=500,80 (APC1001-0047.D)



RECEIVED

APPROVED	O.G. FIG.	
BY	CLASS	SUBCLASS
GRAFTSMAN		

WO 98/25971


09/214251

PCT/GB97/03400

13/14

FIG. 12

(b) HPLC analysis of purified Fab'-PEG.



14/14

O.G. FIG.		
APPROVED	BY	CLASS
DRAFTSMAN		SUBCLASS

Pharmacokinetics of ^{125}I labelled anti-PDGFR IgG, Fab' and FAB'-PEG in rats.




FIG.13